

First-Order Signals in Compact QED with Monopole Suppressed Boundaries[†]

Th. Lippert^a, A. Bode^a, V. Bornyakov^b, and K. Schilling^{a,c}

^aDepartment of Physics, University of Wuppertal, D-42097 Wuppertal, Germany

^bInstitute of High Energy Physics, 142284 Protvino, Russia

^cHLRZ c/0 KFA, D-52425 Jülich, Germany

Pure gauge compact QED on hypercubic lattices is considered with periodically closed monopole currents suppressed. We compute observables on sublattices which are nested around the centre of the lattice in order to locate regions where translation symmetry is approximately recovered. Our Monte Carlo simulations on 24^4 -lattices give indications for a first-order nature of the U(1) phase transition.

1. INTRODUCTION

The presence of metastabilities at the phase transition of compact QED became obvious in simulations on 6^4 -lattices with periodic boundary conditions (p.b.c.)[1]. With increasing lattice size, there was growing confidence that the U(1) phase transition is exhibiting a weakly first-order nature, culminating in the observation of extremely strong hysteresis effects appearing in Monte Carlo series on 16^4 -lattices[2].

This widely accepted opinion has been questioned recently. The authors of Refs. [3,4] have implemented compact U(1) on a lattice with trivial homotopy group, *i.e.* SH_4 , the surface of H_5 . No indications for metastabilities have been found at β_c in simulations on lattices of different sizes—up to a number of degrees of freedom equivalent to a 16^4 -lattice with hyper-toroidal topology. Similarly, fixed boundary conditions (f.b.c.) hint at a continuous transition[5,6].

The implementation of U(1) on SH_4 as well as with f.b.c. explicitly breaks the translation symmetry and produces inhomogeneities of the free energy. One might argue that the absence of first-order effects can be explained in these terms. Alternatively it has been suggested in Ref. [4] that the existence of topologically non-

trivial monopole loops wrapping around the lattice due to p.b.c. might lead to a first-order signature at β_c .

We can check this assumption by studying the model with p.b.c. but periodically closed monopole loops forbidden and looking for first-order signals. For this purpose, we suppress monopole currents emerging in the Monte Carlo update that would point outside the boundary shell and thus would be closed due to p.b.c.

As monitoring quantity we measure the average plaquette on the hyper-surface of subcubes nested around the centre of the lattice and we try to establish a region where translation symmetry is recovered approximately.

2. SUPPRESSION OF MONOPOLES ON BOUNDARY

We suppress monopoles on the boundaries in such a way that spinwaves can penetrate them. The monopole current is defined by[7]

$$\begin{aligned} m_\mu(x) &= \frac{1}{2} \epsilon_{\mu\nu\rho\sigma} [n_{\rho\sigma}(x+\nu) - n_{\rho\sigma}(x)], \\ n_{\rho\sigma} &= -2, -1, 0, 1, 2, \end{aligned} \quad (1)$$

where the number of Dirac sheets $n_{\mu\nu}(x)$ is given by the difference

$$\frac{1}{2\pi} [\bar{\Phi}_{\mu\nu}(x) - \Phi_{\mu\nu}(x)] \quad (2)$$

[†]Work supported by the Deutsche Forschungsgemeinschaft under grant No. Schi 257/3-2, Schi 257/1-4 and by EC contract CHRX-CT92-0051.




Figure 1. Plaquettes contributing to $m_\mu(x)$.




Figure 2. Measurements on nested shells.

between physical flux $\bar{\Phi}_{\mu\nu}(x)$ and plaquette flux $\Phi_{\mu\nu}(x)$ [7]. For a link U_ρ , to be updated in a boundary-cube with normalvector e_μ , we search for the plaquettes associated with the Dirac sheets $n_{\rho\sigma}(x)$ that contribute to the monopole current $m_\mu(x)$ being parallel to e_μ . These plaquettes are depicted in Fig. 1 with dotted lines. If, for any of the corresponding cubes, the change in flux is larger than 2π , the proposed link change is rejected in the Monte Carlo. This particular update rule fulfills *detailed balance* and is equivalent to an infinitely large chemical potential for the monopoles on the boundary shell[8] with the action $S = S_W + S_\lambda$. S_W is the standard Wilson action and S_λ is the additional term which suppresses monopole currents m_μ at the lattice slice $x_\mu = 0$:

$$S_\lambda = \lambda \cdot \sum_\mu \sum_{x: x_\mu=0} |m_\mu(x)|, \quad \lambda \rightarrow \infty. \quad (3)$$

S_λ explicitly breaks translation symmetry.

3. RESULTS

On the 16^4 - as well as on the 24^4 -lattice, we first scanned the β -range in order to get a rough idea about the location of the (pseudo)-critical region. It is shifted towards smaller β -values as expected from monopole suppression. Table 1 shows the number of Metropolis sweeps we performed subsequently for the such chosen β -values and the plaquette averaged over the whole lattice. On the 24^4 -lattice in the average about 1% of the proposed updates are rejected at the boundary.

The plaquettes have been sampled on the surfaces of sub-hypercubes nested around the centre




Figure 3. Average plaquette (dotted line) and plaquettes on shells 1,3,5 and 7 (from top to bottom) on the 16^4 -lattice.

of the lattice. The 3D projection of these shells is shown in Fig. 2. The boundary shell is denoted by the index 1, the inner shell gets the index $L/2$.

Table 1: Lattice sizes, β -values and # of sweeps.

size	β	# of sweeps	$\langle P \rangle$
16^4	0.990	30000	0.6290(2)
	0.995	10000	0.6408(4)
	1.000	60000	0.6530(2)
	1.002	60000	0.6584(1)
	1.003	60000	0.6599(1)
	1.005	60000	0.6630(1)
	1.010	10000	0.6688(2)
24^4	1.0060	60000	0.6546(1)
	1.0065	60000	0.6572(1)
	1.0070	250000	0.65856(5)
	1.007125	200000	0.65935(9)
	1.0075	80000	0.66010(9)
	1.0078	50000	0.66099(6)
	1.0080	55000	0.66139(5)

The boundary shell is always in a deconfined state. This is a well known phenomenon in the absence of monopoles[8]. Therefore, the variation of the plaquette value within the metastable region is very small for this shell. As the boundary shell contains the outweighing part of the sites, the plaquette average over the whole lattice is dominated by the contribution of the boundary. We found the influence of the boundary decreasing towards the centre, *i.e.* the plaquette values show increasing ‘steps’ (as functions of β) going to the inner shells, see Fig. 3 for the 16^4 -lattice. For inner shells the plaquette values are close to




Figure 4. Time series on 24^4 -lattice for different shells.

each other and their behaviour is more and more like that of the plaquette of the H_4 system with p.b.c. We interpret this results as the onset of the recovery of translation invariance around the centre.

We carefully analyzed the time series for the inner shells on the 16^4 -lattice but the series merely exhibited a chaotic behaviour and no metastabilities could be identified. Therefore we decided to work on a very large 24^4 -system and we performed altogether nearly one million sweeps.

On this lattice, we found a much broader region in 4-d space at the centre where the behaviour of the system resembled that of $U(1)$ on H_4 with p.b.c. and we found evidence for metastabilities in the time series. In Fig. 4, the time series of plaquettes for the shells 1, 5 and 9 are depicted for $\beta = 1.007125$. At this value of β we encountered long metastabilities extending over more than 10000 sweeps as can be most clearly seen in the series of shell 9. As the fluctuations of the plaquette in inner shells are larger than the gap between the two metastable states we performed a windowing procedure in order to average out the fluctuations on small scales and applied the method to the time series of shell 9. The result is presented in Fig. 5, where we see a non-ambiguous double-peak structure of the histogram. More detailed results will be presented in a forthcoming




Figure 5. Histogram of shell 9 after removing of fluctuations.

paper.

4. CONCLUSIONS

In our simulations

- we did not observe first order signals on small lattices because of boundary effects. We believe this is true for f.b.c. as well.
- we observed metastabilities coming up on the 24^4 -lattice. This implies that periodically closed monopole loops can not be the reason for the first order nature of $U(1)$ with p.b.c.

REFERENCES

1. J. Jersák, T. Neuhaus and P. M. Zerwas, Phys. Lett. No. 133B (1983) 103.
2. see e.g. Th. Lippert, G. Bhanot, K. Schilling and P. Ueberholz, Nucl. Phys. B (Proc. Suppl.) 30 (1993) 912.
3. Ch. Lang and T. Neuhaus, Nucl. Phys. B (Proc. Suppl.) 34 (1994) 543.
4. Ch. Lang and T. Neuhaus, Preprint Bielefeld University, 1994.
5. M. Baig and H. Fort, UAB-FT-338, 1994.
6. K.-H. Mütter and K. Schilling, Nucl. Phys. B 200 (1982) 362.
7. T. DeGrand and D. Toussaint, Phys. Rev. D22 (1980) 2478.
8. V. Bornyakov, V. Mitrjushkin and M. Müller-Preussker, Nucl. Phys. B (Proc. Suppl.) 30 (1993) 587.